

Neogra 100mg Oral Jelly

Neogra Oral Jelly (Sildenafil Citrate Jelly)

Product Information: Neogra Jelly is an oral medication for the treatment of erectile dysfunction. The active ingredient contained in Neogra oral jelly is Sildenafil Citrate . It starts working about 30 minutes after taking the dose and the medical effect will last for 3 to 4 hours. Each 100 mg sachet of Neogra Oral Jelly contains 100 mg of the active ingredient Sildenafil Citrate .

Active Ingredient present in Neogra Jelly: The active ingredient present in Neogra Jelly is Sildenafil Citrate.

Uses of Sildenafil Oral Jelly: Neogra jelly is an oral gel solution of Sildenafil and it is use for the treatment of erectile dysfunction in men. Neogra Jelly contains 100mg Sildenafil Citrate - the exact same active ingredient as in Sildenafil tablets or Generic Sildenafil tablets. Neogra Jelly is supplied in a box containing 7 different flavors. The flavor combinations are either pineapple, orange, strawberry, vanilla, banana, black currant, and butterscotch or mint, rose, mango, lemon, cherry, chocolate, and watermelon. It is not possible to specify a specific flavor combination upon ordering Neogra Jelly.

Neogra Oral Jelly Dosage: Use one sachet of Neogra Oral jelly as and when required, approximately half an hour to one hour before sexual activity. The maximum recommended dosing frequency is one sachet of Neogra Oral Jelly (Neogra gel) per day. In other words, you can take another dose only if 24 hours have passed since your last dose.

How to Use Neogra Oral Jelly: Neogra Oral Jelly comes in sachets of 100mg Sildenafil Citrate each. Cut open the sachet and squeeze the jelly into the mouth and swallow the entire content of the sachet. The Neogra Oral Jelly solution melts very fast and is absorbed by the body's enzymes in approximately 30 min after taking the medication. Neogra Jelly works for about 2-3 hours after consumption and is thus known as the quick and fun solution to an instant erection.

Neogra 100mg Oral Jelly

Side Effects of Neogra Oral Jelly: Neogra Jelly has similar side effects as other Generic Sildenafil Citrate medications. These are usually mild to moderate and usually don't last longer than a few hours. Some of these side effects are more likely to occur with higher doses. The most common side effects of Neogra Jelly are headache, flushing of the face, and upset stomach.

Less common side effects that may occur are temporary changes in color vision (such as trouble telling the difference between blue and green objects or having a blue color tinge to them), eyes being more sensitive to light, or blurred vision.

In rare instances, men taking PDE5 inhibitors (oral erectile dysfunction medicines, including this medicine) reported a sudden decrease or loss of vision in one or both eyes. It is not possible to determine whether these events are related directly to these medicines, to other factors such as high blood pressure or diabetes, or to a combination of these. If you experience sudden decrease or loss of vision, stop taking PDE5 inhibitors, including Neogra and call a doctor right away.

In rare instances, men have reported an erection that lasts many hours. You should call a doctor immediately if you ever have an erection that lasts more than 4 hours. If not treated right away, permanent damage to your penis could occur.

Sudden loss or decrease in hearing, sometimes with ringing in the ears and dizziness, has been rarely reported in people taking PDE5 inhibitors, including Neogra Jelly. It is not possible to determine whether these events are related directly to the PDE5 inhibitors, to other diseases or medications, to other factors, or to a combination of factors. If you experience these symptoms, stop taking Neogra Jelly and contact a doctor right away.

Heart attack, stroke, irregular heartbeats, and death have been reported rarely in men taking Neogra . Most, but not all, of these men had heart problems before taking Neogra. It is not possible to determine whether these events were directly related to Neogra Oral Jelly .

If you want more information or develop any side effects or symptoms you are concerned about, call your doctor. In case of accidental overdose, call your doctor right away.

Neogra 100mg Oral Jelly

Warnings & Precautions When Using Neogra Oral Jelly: Do not consume more than one sachet of Neogra Jelly (Oral gel solution) per 24 hours. Some conditions and medicines interfere with the natural erection process. The penis cannot fill with enough blood. The man cannot have an erection. This is called erectile dysfunction if it becomes a frequent problem. During sex, your heart works harder. Therefore sexual activity may not be advisable for people who have heart problems. Before you start any treatment for erectile dysfunction, ask your doctor if your heart is healthy enough to handle the extra strain of having sex. If you have chest pains, dizziness or nausea during sex, stop having sex and immediately tell your doctor you have had this problem. Neogra Jelly can cause mild, temporary lowering of your blood pressure. You will need to have a thorough medical exam to diagnose your erectile dysfunction and to find out if you can safely take Neogra alone or with your other medicines.

Be sure to tell your doctor if you:

- have ever had any heart problems (e.g., angina, chest pain, heart failure, irregular heartbeats, heart attack or narrowing of the aortic valve)
- have ever had a stroke.
- have low or high blood pressure.
- have ever had severe vision loss.
- have a rare inherited eye disease called retinitis pigmentosa.
- have ever had any kidney problems.
- have ever had any liver problems.
- have ever had any blood problems, including sickle cell anemia or leukemia.
- are allergic to sildenafil (the active ingredient).
- have a deformed penis, Peyronie's disease, or ever had an erection that lasted more than 4 hours.
- have stomach ulcers or any types of bleeding problems & are taking any other medicines.

Neogra 100mg Oral Jelly

Tell your doctor about any medicines you are taking. Do not start or stop taking any medicines before checking with your doctor. Neogra should never be used with medicines that contain nitrates. If you are taking medicines called alpha-blockers for the treatment of high blood pressure or prostate problems, your blood pressure could suddenly drop. You could get dizzy or faint. If you are taking a protease inhibitor, your dose may be adjusted.

Neogra Jelly should not be used with any other medical treatments that cause erections. These treatments include pills, medicines that are injected or inserted into the penis, implants or vacuum pumps.

Neogra Jelly only for patients with erectile dysfunction. It is not for newborns, children, or women. Neogra Jelly not protect you or your partner from getting sexually transmitted diseases, including HIV—the virus that causes AIDS. It is not a hormone or an aphrodisiac.

Disclaimer: The above information is provided to the best of our knowledge and in good faith, it is without a warrant of any kind, expressed or implied.